



A novel technique for invasive aortic valve pressure gradient measurement using a 6 Fr Swan-Ganz catheter: a case series

Nayan Agarwal¹, Ujjwal Rastogi¹, Ahmed N. Mahmoud ², and Deepak L. Bhatt ^{3*}

¹Department of Cardiology, Cardiovascular Institute of the South, 225 Dunn Street, Houma, LA 70360, USA; ²Department of Medicine, University Hospital Portage Medical Center, 6847 N Chestnut St Ravenna, OH 44266, USA; and ³Department of Medicine, Brigham and Women's Hospital Heart & Vascular Center, Harvard Medical School, 75 Francis St, Boston, MA 02115, USA

Received 8 April 2021; first decision 30 April 2021; accepted 14 September 2021; online publish-ahead-of-print 24 September 2021

Background

Simultaneous left ventricular (LV) and aortic (Ao) pressure gradient assessment has been rendered challenging since the recall of the Langston catheter. Here we describe a simple method for simultaneous LV and Ao pressure gradient assessment using a Swan-Ganz catheter.

Case summary

We describe two cases where assessment of simultaneous left ventricle and Ao valve gradients was done using a Swan-Ganz catheter to assess the degree of Ao stenosis and dynamic LV outflow obstruction.

Discussion

Using Swan-Ganz catheter assessment of simultaneous left ventricle and Ao valve gradients can simplify the procedure with reduced cost and increased patient safety.

Keywords

Case series • Dual lumen catheter • Aortic stenosis • Swan-Ganz

Learning points

- The Swan-Ganz catheter is a viable alternative for simultaneous assessment of the left ventricle–aortic (Ao) pressure gradient without a dual lumen Langston catheter.
- Using the Swan-Ganz catheter for Ao valve studies can reduce procedural cost and complications.

Introduction

Accurate assessment of aortic (Ao) stenosis when there is discrepancy between echocardiographic and clinical data requires

invasive assessment of simultaneous left ventricular (LV) and Ao pressure gradients using a dual lumen catheter.¹ Due to the recall of the Langston dual lumen catheter,² such haemodynamic assessment has been rendered challenging. Various techniques have been described by operators and shared on social media to obtain LV–Ao gradients such as using a mother in child technique with a 6 or 7 Fr guide, a long sheath with a 4 Fr catheter inside, a fractional flow reserve (FFR) wire extending from the catheter, or dual arterial access.³ All these techniques can be associated with increased cost, time, equipment, and complications from the larger arteriotomy size and need for dual arterial access.^{4,5} Here we describe a simple and cost-effective technique of obtaining simultaneous LV–Ao gradients.

* Corresponding author. Tel: 1-857-307-4071, Email: dlbhattmd@post.harvard.edu

Handling Editor: Luigi Biasco

Peer-reviewers: Romain Didier and Dejan Milasinovic

Compliance Editor: Daniel Tardo

Supplementary Material Editor: Nida Ahmed

© The Author(s) 2021. Published by Oxford University Press on behalf of the European Society of Cardiology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

Timeline

Case 1

Day 1	Clinical assessment—shortness of breath
Day 4	Outpatient echocardiogram suggestive of mild–moderate aortic (Ao) stenosis
Day 10	Outpatient heart catheterization with the described technique to assess the degree of Ao stenosis

Case 2

Day 1	Clinical assessment—shortness of breath
Day 3	Outpatient echocardiogram suggestive of left ventricular (LV) outflow tract obstruction
Day 7	Outpatient heart catheterization with the described technique to assess the degree of LV outflow tract obstruction




Figure 1 Swan-Ganz catheter in left ventricle. Blue arrow, distal transducer port position; red arrow, proximal transducer port position.

Case presentation

Through the slender 5/6 Fr sheath placed in the radial artery, the Ao valve was crossed using a Judkins Right 4 (JR4) catheter and 0.035" straight tip wire. The catheter was introduced in the LV over this wire which was then swapped out for 300 cm 0.014" Terumo Runthrough wire over which the JR4 catheter was exchanged for a 6 Fr Swan-Ganz catheter (Edwards), which was used earlier for the right heart catheterization, with the distal port in the LV and the proximal port in the Ao ([Figure 1](#)). The balloon on the Swan-Ganz catheter was deflated while the catheter was being advanced, and once in the LV, the balloon was inflated to help move the catheter away from LV wall, which might have caused pressure waveform dampening. Thus, simultaneous pressures could be obtained for the invasive haemodynamic assessment of Ao stenosis ([Figure 2](#)). Since the Swan-Ganz catheter is an end-hole catheter, this technique can also be used for the assessment of dynamic LV outflow obstruction ([Figure 3](#)), without the need for dual arterial access. In cases of subclavian tortuosity, the exchange can also be done over a long 0.025" Swan wire or a 0.018" wire, which might be easier.

Case 1

A 76-year-old male with history of hypertension and hyperlipidaemia presented with worsening shortness of breath. Cardiac workup showed a moderate risk stress test and mild–moderate Ao stenosis on echocardiogram with a mean gradient of 18 mmHg, Ao valve area of 1.4 mm², and peak velocity of 3.06 m/s ([Figure 4](#)). Due to body habitus, the echocardiogram images were not diagnostic. He also had an unremarkable chest X-ray and pulmonary function testing. Due to ongoing symptoms, the decision was made to proceed with heart catheterization which showed mild non-obstructive coronary artery disease and normal filling pressures. Mild Ao stenosis was confirmed with simultaneous LV–Ao gradients assessed via the above-described technique ([Figure 2](#)).

Case 2

A 51-year-old female with no cardiac risk factors presented with progressively worsening shortness of breath. Physical exam was consistent with a systolic murmur which increased with vagal manoeuvres. Echocardiogram showed normal ejection fraction, normal valvular structures with asymmetric septal hypertrophy, and turbulence in the LV outflow tract with a peak gradient of 56 mmHg at rest ([Figure 5](#)). The stress test was high risk. Heart catheterization showed normal coronary arteries and normal filling pressures. Simultaneous LV–Ao pressure gradients performed with the above technique using a Swan-Ganz catheter showed a resting peak LV outflow tract gradient of 113 mmHg, which increased to 164 mmHg with induction of premature ventricular contractions. The pullback confirmed that the obstruction was in the LV outflow tract ([Figure 3](#)).

Discussion

The above-described technique can be very useful in haemodynamic assessment of Ao stenosis or LV outflow tract obstruction in cases where the echocardiographic assessment may not be accurate and invasive gradient measurements are needed. With this technique, no extra equipment, larger or additional arteriotomy was needed. Therefore, this approach can help reduce costs and complications. Since the Swan-Ganz is an end-hole catheter, there can be pressure dampening if the catheter is against the LV wall. This can be overcome by balloon inflation and gentle readjustment of catheter position. To avoid the risk of air embolization from balloon rupture during balloon inflation, we recommend using saline or contrast for the balloon inflation. To the best of our knowledge, our report was the first to describe the above technique.³

The gradients obtained by this technique in the two cases above closely matched the echo gradients, and the waveforms appear undamped, which makes us believe that the assessment of gradients is very accurate. In the future, if the results of this method can be




Figure 2 Simultaneous left ventricular and aortic gradient measurement across a stenosed aortic valve. Ao, aortic; Blue tracing, left ventricle; D, diastole; EDP, end-diastolic pressure; LV, left ventricular; Red tracing, aorta; S, systole.

validated by using two separate end-hole catheters simultaneously in the LV and Ao, then this technique can completely replace the need for a Langston dual lumen catheter for such assessment.

Lead author biography



Dr Nayan Agarwal is an interventional cardiologist at Cardiovascular Institute of the South in Louisiana. He finished his medical school in 2009 at Maulana Azad medical college in India and then pursued internal medicine residency followed by fellowship in cardiology and interventional cardiology at University of Florida, Gainesville.

Supplementary material

Supplementary material is available at *European Heart Journal - Case Reports* online.

Slide sets: A fully edited slide set detailing this case and suitable for local presentation is available online as [Supplementary data](#).

Consent: The authors confirm that written consent for submission and publication of this case report including images and associated text has been obtained from the patient in line with COPE guidance.

Conflict of interest: D.L.B. discloses the following relationships—Advisory Board: Cardax, CellProthera, Cereno Scientific, Elsevier Practice Update Cardiology, Janssen, Level Ex, Medscape Cardiology, MyoKardia, Novo Nordisk, PhaseBio, PLx Pharma, Regado Biosciences; Board of Directors: Boston VA Research Institute, Society of Cardiovascular Patient Care, TobeSoft; Chair: American Heart Association Quality Oversight Committee; Data Monitoring Committees: Baim Institute for Clinical Research (formerly Harvard




Figure 3 Assessment of dynamic left ventricular outflow obstruction. (A) Resting gradient between left ventricle and aorta. (B) Increasing gradient between left ventricle and aorta with induction of premature ventricular contraction with reduction in pulse pressure of aorta. (C) Pullback showing loss of left ventricle and aortic pressure gradient when catheter pulled back into left ventricular outflow tract. Blue tracing, left ventricle; D, diastole; EDP, end-diastolic pressure; Red tracing, aorta; S, systole.

Clinical Research Institute, for the PORTICO trial, funded by St. Jude Medical, now Abbott), Cleveland Clinic (including for the ExCEED trial, funded by Edwards), Contego Medical (Chair, PERFORMANCE 2), Duke Clinical Research Institute, Mayo Clinic, Mount Sinai School of Medicine (for the ENVISAGE trial, funded by Daiichi Sankyo), Population Health Research Institute; Honoraria: American College of Cardiology (Senior Associate Editor, Clinical Trials and News, ACC.org; Vice-Chair, ACC Accreditation Committee), Bain Institute for Clinical Research (formerly Harvard Clinical Research Institute; RE-DUAL PCI clinical trial steering committee funded by Boehringer Ingelheim; AEGIS-II executive committee funded by CSL Behring), Belvoir Publications (Editor in Chief, Harvard Heart Letter), Canadian Medical and Surgical Knowledge Translation Research Group (clinical trial steering committees), Duke Clinical Research Institute (clinical trial steering committees, including for the PRONOUNCE trial, funded by Ferring Pharmaceuticals), HMP Global (Editor in Chief, Journal of Invasive Cardiology), Journal of the American College of Cardiology (Guest Editor; Associate Editor), K2P (Co-Chair, interdisciplinary curriculum), Level Ex,

Medelligence/ReachMD (CME steering committees), MJH Life Sciences, Population Health Research Institute (for the COMPASS operations committee, publications committee, steering committee, and USA national co-leader, funded by Bayer), Slack Publications (Chief Medical Editor, Cardiology Today's Intervention), Society of Cardiovascular Patient Care (Secretary/Treasurer), WebMD (CME steering committees); Other: Clinical Cardiology (Deputy Editor), NCDR-ACTION Registry Steering Committee (Chair), VA CART Research and Publications Committee (Chair); Research Funding: Abbott, Afimmune, Amarin, Amgen, AstraZeneca, Bayer, Boehringer Ingelheim, Bristol-Myers Squibb, Cardax, CellProthera, Cereno Scientific, Chiesi, CSL Behring, Eisai, Ethicon, Ferring Pharmaceuticals, Forest Laboratories, Fractyl, Garmin, HLS Therapeutics, Idorsia, Ironwood, Ischemix, Janssen, Lexicon, Lilly, Medtronic, MyoKardia, Novartis, Novo Nordisk, Owkin, Pfizer, PhaseBio, PLx Pharma, Regeneron, Roche, Sanofi, Synaptic, The Medicines Company, 89Bio; Royalties: Elsevier (Editor, Cardiovascular Intervention: A Companion to Braunwald's Heart Disease); Site Co-Investigator: Abbott, Biotronik, Boston Scientific, CSI, St. Jude Medical (now



Figure 4 Echocardiogram continuous wave through aortic valve.




Figure 5 Echocardiographic resting gradient through left ventricular outflow tract.

Abbott), Svelte; Trustee: American College of Cardiology; Unfunded Research: FlowCo, Merck, Takeda. All other authors declared no conflict of interest.

Funding: None declared.

References

1. Saikrishnan N, Kumar G, Sawaya F, Lerakis S, Yoganathan A. Accurate assessment of aortic stenosis: a review of diagnostic modalities and hemodynamics. *Circulation* 2014;**129**:244–253.
2. U.S. Food & Drug Administration. Vascular Solutions, Inc. Recalls Langston Dual Lumen Catheter Due to Risk of Separation During Use. <https://www.fda.gov/medical-devices/medical-device-recalls/vascular-solutions-inc-recalls-langston-dual-lumen-catheter-due-risk-separation-during-use> (21 July 2021).
3. @nayanagarwal85. No Langston. LV-Ao Done with 6 Fr Swan Which Is Already on Table for the RHC. No Need for Extra Equipment or 7 Fr or Dual Access. Posted 11 February 2021. <https://t.co/TSwSuBtdDs> (21 July 2021).
4. Garg J, Tyagi S, Allana S. Assessing transvalvular aortic gradient with Swan-Ganz catheter. *JACC Cardiovasc Interv* 2021;**14**:e123–e124.
5. Bangalore S, Bhatt DL. Images in cardiovascular medicine. Right heart catheterization, coronary angiography, and percutaneous coronary intervention. *Circulation* 2011;**124**:e428–e433.